

January 7, 2011

VACCINATION PROTOCOL

Calves 5 weeks of age – ear punch for BVD prior to vaccinations

- IBR, BVD, P13, BRSV (modified live)
- 5-way Lepto
- 7-way Clost

Bova Shield

Ultrabac 7/ Somubac

Calves 3-4 weeks after initial immunizations

- 7-way Clostridium group, HS
- IBR, BVD, P13, BRSV (modified live)
- 5-way Lepto

Ultrabac 7/ Somubac

Bova Shield

Calves 6 months of age

- IBR, BVD, P13, BRSV (modified live)
- 5-way Lepto

Bova Shield

Calves 12 months of age (1-2 months prior to breeding)

- IBR, BVD, P13, BRSV (all modified live)
- 5-way Lepto, 7-way clostridium, HS

Bova Shield

Ultrabac 7/ Somubac

Heifers during pregnancy check (16-18 months of age)

- 5-way Lepto *Leptoferm-5*

Pregnant heifers and cows, 6.5 months gestation

- J-5 (#1/5)

J-5 Bacterin

Pregnant heifers and cows, 7 months gestation (dry-off)

- J-5 (#2/5)
- 7-way Clostridium group, HS

J-5 Bacterin

Ultrabac 7/ Somubac

Pregnant heifers and cows 7.5 months gestation

- K99, Rota and Coronaviruses (#1/2)

Scourguard 3 (K)/C

Pregnant heifers and cows 8 months gestation

- J-5 (#3/5)

J-5 Bacterin

Pregnant heifers and cows 8.5 months gestation or 7-10 days after previous vaccination

- K99, Rota and Coronaviruses (#2/2)

Scourguard 3 (K)/C

Lactating heifers (1st parity) and cows 28 days in milk

- J-5 (#4/5)

J-5 Bacterin

Lactating heifers and cows 20-35 days in milk

- IBR, BVD, P13, BRSV (all modified live)
- 5-way Lepto

Bova Shield

Lactating heifers and cows 90-110 days in milk

- J-5 (#5/5)

J-5 Bacterin

Lactating heifers and cows during pregnancy check

- 5-way Lepto

Leptoferm 5

Any animal that is 340 days since last MLV will get boosted with MLV

Bova Shield

Rabies (*Defensor*) should be given in late winter or early spring, 4 to 6 weeks prior to the expected date that animals will be turned-out to pasture. Rabies should be given to all members of the herd older than 4 months of age. Initial dose should consist of a single injection, followed with a booster on an annual basis. Animals not old enough to receive the vaccinations during the spring should be vaccinated during the subsequent summer months when deemed appropriate by the Manager and Clinical Veterinarian.

Full names on abbreviated diseases:

BVD	Bovine viral diarrhea
IBR	Infectious bovine rhinotracheitis
Lepto	Leptospirosis
BRSV	Bovine respiratory syncytial virus
P13	Parainfluenza-3
HS	Hemophilus